Historic, Archive Document

Do not assume content reflects current scientific knowledge, policies, or practices.

| 1 | | | | |
|---|--|--|--|----|
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | 1. |
| | | | | |
| | | | | 4 |
| | | | | |
| | | | | |
| | | | | |
| | | | | * |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | 1 |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |

XXXXXX RETAIL

PLANTSMEN



BROMELIA COLLECTION

U. S. Department of Agriculture

| AECHMEA | | |
|---|----------------------|---------------------------------------|
| Fasciata | 3-4-5" | 1.25 - 2.00 - 3.50 |
| Fulgens | 4-5" | 2,50 - 3,50 |
| Fulgens discolor | 4-5" | 4.00 - 6.00 |
| Speciosa miniata (gray | 411 | 6.00 |
| Tinctoria | 4-511 | 3,50 - 5.00 |
| ANANAS | 1. 0 | |
| Cochinchinensis | 5-711 | 25,00 8:35,00 |
| Porteana | 50 | • |
| | J | 7.50 |
| AREGELIA | 4 5 3 | 0.50 7.50 |
| Marmorata | 4-5 | 2.50 - 3.50 |
| Spectabilis | 4-51 | 2,50 - 3,50 |
| BILLBERGIA | | |
| Alberta | 3-4" | 1,00 - 1,50 |
| Alberta magnifica | 3-4" | 1.25 - 2.00 |
| Forgetti | 5 ¹¹ | 5.00 - 7.50 |
| Hybrid Nutans & Beucke: | ri 3-4" | .75 - 1.25 |
| Leopoldi | 411 | 2.50 |
| Morelli | 3-411 | 1.00 - 1.50 |
| | 3-3-11 | |
| Macrocalyx | 21,4311 | 1,00 - 1,50 |
| Nutans | | .5075 |
| Perringeana | 4-5" | 7.50 - 10.00 |
| Pyramidalis | 3 <u>3</u> " | 1.50 |
| Quintitissima | 3-3 ½ " | ·75 - 1.00 |
| Rhodecyana | 4-51 | 5,00 - 7,50 |
| Sandersii | 3-3-3-4" | 1,25 - 1,75 - 2,50 |
| Zebrina | 411 | 5,00 |
| GUZMANIA | | |
| Lindeni | 311 | 7:50 |
| Musaica | 411 | 10:00 |
| Zahnii | 4-5" | 3,50 - 5,00 |
| | 4-0 | 0,50 - 3,00 |
| NIDULARIUM | 4-511 | F 00 7:50 |
| Amazonica | | 5.00 - 7.50 |
| Innocenti | 4" | 3,50 |
| TILLANDSIA | | |
| Lindeni | | 5 ₆ 00 - 7 ₆ 50 |
| VRIESIA | | , |
| Erecta | 411 | 5,00 |
| Favorita | 4-5" | 3 ,50 - 5,00 |
| Guttata | 311 | 3,50 |
| Hieroglyphlica | 4" 4" 3" | 7.50 |
| Magnifica | <u>, 11</u> | 7,50 |
| | 711 | |
| Splendens major | J | 7.50 |
| CRYPTANTHUS | $2^{\frac{1}{2}}$ 11 | |
| acaulis | | .35 |
| acaulis | 2 <u>‡</u> n 3n | •35 |
| bivittatus | 3" | 1,00 |
| beuckeri | 2111 | ·, 35 ² 75 |
| roseus pictus | 2 ± -3" | 50 - 50 |
| terminalis | 24-3" | |
| zonatus | 2 1 -311 | 75 - 1:00 75 - 1:00 |
| zonatus zebrinus | 3-4-4=" | 1.00 - 1.50 - 2.50 |
| the manner of other than to be the black of the | 6 | |

budado term Enterior 1